

## Ghana Summer School

### I. Melting Ice

Supply List:

- Two Large Containers
- Fresh water
- Salt
- ice cubes
- Food coloring

*Two containers hold equal amounts of water and ice; however, one container is filled with fresh water and one is filled with salty water. In which container will the ice melt the fastest? Why?*

Now let's test your theory. Fill both containers equally with about 1600 ml of fresh water. Mix 35 ml of salt into container #2 (make sure all the salt is dissolved). Place a few ice cubes in each container. Observe what happens.

*Now drop a (very) little food coloring on to the ice in the containers. What does this reveal? Once the ice has completely melted in one of the containers, discuss how you might induce the ice to melt faster in the other container. Sketch a profile of temperature, salinity, and density in each cup at the beginning, in the middle, and at the end of the experiment.*

## II. Coriolis Machine

### Supply List:

Plastic Sheet

Paper with Reference Frame

Thumbtack

Dry Erase Marker

The Coriolis Machine is composed of a plastic sheet placed over a white piece of paper secured together in the center with the thumbtack. The plastic sheet is your rotating reference (non-inertial) frame, as it can be easily rotated over the white paper (which represents the non-rotating or inertial reference frame). To measure a rotation rate, radial lines of equal spacing are drawn through the center of the paper. The distance between radial lines marks the rotation angle over one time step  $\Delta t$ . Paths of particles moving in straight lines are also marked on the inertial reference frame. Ticks on the paths mark the distance the particle traveled in one time step. *Considering the spacing of the particles, can you identify which line represents an accelerating particle?*




Fig. 1. Exploded view of Coriolis machine.

Now you are ready to run the machine. Start by marking the initial position of the particles with a small x. Also mark one of the radial lines as a reference for turning. Now rotate the non-inertial reference frame counter-clockwise one time step, i.e., move your reference radial line over one position to the right. Now mark the new location of the particle by placing another mark at the second tick on the particle path. Continue to rotate and mark the particle positions until you understand the path of the particles on the non-inertial (plastic) reference frame.

*Which direction are the particles deflecting? Does one particle deflect more than the other particle? What happens if you try the machine again, rotating clockwise? Try it!*